



SUBDIFFERENTIAL DETERMINATION OF A PRIMAL LOWER REGULAR FUNCTION ON A BANACH SPACE

Milen Ivanov¹, Matey Konstantinov^{2,*}, Nadia Zlateva³

¹Radian Life Technologies Ltd., Nicosia, Cyprus, e-mail:milen@radian-life.com

²Faculty of Mathematics and Informatics, Sofia University, 5, James Bourchier Blvd, 1164 Sofia, Bulgaria,
e-mail:matey@fmi.uni-sofia.bg

³Faculty of Mathematics and Informatics, Sofia University, 5, James Bourchier Blvd, 1164 Sofia, Bulgaria,
e-mail:zlateva@fmi.uni-sofia.bg

ABSTRACT

We generalise to Banach space Thibault-Zagrodny Theorem that if f and g are primal lower regular and $\partial f = \partial g$ locally, then f and g locally differ by a constant. Our method consists in reduction to slopes considerations.

Keywords primal lower regular function · slope · determination of a function

References

- [1] M. Ivanov, N. Zlateva: *Slopes and Moreau-Rockafellar theorem*, J. Convex Analysis 32/2 (2025) 359–374.
- [2] R. A. Poliquin: *Integration of subdifferentials of nonconvex functions*, Nonlinear Anal. Th. Meth. Appl. 17 (1991) 385–398
- [3] L. Thibault, *Unilateral Variational Analysis in Banach Spaces. Part I: General Theory. Part II: Special Classes of Functions and Sets*, World Scientific (2023) ISBN: 978-981-125-816-9.

*Corresponding Author's E-mail: matey@fmi.uni-sofia.bg